

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/00	A2	(11) International Publication Number: WO 99/03446 (43) International Publication Date: 28 January 1999 (28.01.99)
<p>(21) International Application Number: PCT/US98/14666</p> <p>(22) International Filing Date: 15 July 1998 (15.07.98)</p> <p>(30) Priority Data: 08/896,194 17 July 1997 (17.07.97) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/896,194 (CIP) Filed on 17 July 1997 (17.07.97)</p> <p>(71) Applicant (for all designated States except US): E-L MANAGEMENT CORP. [US/US]; 767 Fifth Avenue, New York, NY 10153 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SCHNITTGER, Steven, F. [US/US]; 10 Willets Place, Huntington Station, NY 11746 (US). DECLERCQ, Lieve [BE/BE]; Poorthoflaan 77, B-2180 Ekeren (BE).</p> <p>(74) Agent: TSEVDOS, Estelle, J.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
(54) Title: NOVEL ANTIMICROBIAL COSMETIC COMPOSITIONS		
<p>(57) Abstract</p> <p>The present invention relates to cosmetic or pharmaceutical formulations for topical application which comprises an antimicrobial effective amount of <u>3-furan carboxylic acid</u>. The formulation also optionally comprises an anti-irritant effective amount of ferulic acid and an antimicrobial effective amount of <u>2-furan carboxylic acid</u>. The invention also provides cosmetic or pharmaceutical compositions useful in the treatment of inflammation, comprising effective amounts of 2-furan carboxylic acid or 3-furan carboxylic acid, or mixtures thereof.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NB	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL ANTIMICROBIAL COSMETIC COMPOSITIONS

Related Applications

This application is a continuation in part of copending US Serial
5 No. 08/896,194, filed July 17, 1997, the contents of which are
incorporated herein by reference.

Field of the Invention

The invention relates to novel cosmetic compositions. In
10 particular, the invention relates to cosmetic compositions having
unique antimicrobial and antiirritant properties.

Background of the Invention

Water miscible cosmetic and toiletry formulations are
15 frequently susceptible to microbial contamination, due to their
high water content and the nature of ingredients that they often
contain. This is particularly true with the trend toward
"natural" cosmetic ingredients, which are more likely to be
susceptible to contamination than synthetic ingredients may be.
20 The occurrence of microbial contamination in a cosmetic
formulation can result in an unpleasant odor or the
destabilization of an emulsion; this in turn can lead to the
necessity of reformulation or recall of a commercial product.

To counteract this problem, it is often necessary to add
25 antimicrobial chemicals, such as preservatives or biocides, to
the formulation to prevent the growth of microbes that may be
introduced during the manufacturing, filling or use of the
product. However, such additives have recently fallen into
disfavor, in large part because many preservatives are perceived
30 as causing irritation and consumers are now demanding
preservative-free products. Therefore, the market is shifting
toward lower levels of conventional preservatives, and also the
replacement of conventional preservatives with new molecules.

Thus, there is now a strong demand for cosmetic formulations which are non-irritating and free of traditional preservatives, but which will remain stable and free of contamination in the hands of the consumer. The present invention provides such
5 formulations.

Summary of the Invention

The present invention relates to a cosmetic or pharmaceutical formulation for topical application which
10 comprises an antimicrobial effective amount of 3-furan carboxylic acid. In a preferred embodiment, the formulation also comprises an anti-irritant effective amount of ferulic acid, or a derivative thereof. In a particularly preferred embodiment, the formulation also comprises an antimicrobial effective amount of
15 2-furan carboxylic acid. All such formulations are capable of being made free of preservatives, and achieve the desired effects by the use of naturally occurring materials.

The invention also relates to method for reducing the irritant effect of a cosmetic or pharmaceutical formulation
20 comprising adding to the formulation an anti-irritant effective amount of ferulic acid or a derivative or analog thereof, as well as reducing or preventing irritation on the skin by topical application to the skin of a formulation containing ferulic acid or a derivative thereof. The invention further relates to a
25 method for reducing or preventing inflammation which comprises applying to the skin an anti-inflammatory effective amount of 2-furan carboxylic acid, 3-furan carboxylic acid, or a combination thereof. Also encompassed is a method for inhibiting microbial growth in a cosmetic or pharmaceutical formulation which
30 comprises adding to the formulation an antimicrobial effective amount of 3-furan carboxylic acid.

Also provided by the invention are topical cosmetic and pharmaceutical compositions useful in the treatment of

inflammation in the skin comprising effective amounts of 2-furan carboxylic acid, 3-furan carboxylic acid, or a mixture thereof, as well as methods of treating inflammatory reactions in the skin comprising applying such compositions to the skin.

5

Brief Description of the Drawings

Figure 1 illustrates the inhibitory effect of (A) ferulic acid; (B) 2-furan carboxylic(2-furoic) acid; and (C) 3-furan
10 carboxylic(3-furoic) acid on phospholipase A₂ activity.

Figure 2 illustrates the inhibitory effect of 2-furan carboxylic acid(2F) and 3-furan carboxylic acid (3F) on the inhibition of lymphocyte adhesion to endothelial cells.

15

Detailed Description of the Invention

Furan carboxylic acids are naturally occurring degradation products from lignin and cellulose. 2-furan carboxylic acid has previously been disclosed as having antimicrobial activity, and
20 has been said to be particularly effective against mycobacteria(US Pat. No. 5,387,605). However, antimicrobial activity for 3-furan carboxylic acid has not previously been reported; in the context of the present invention, it has now been unexpectedly discovered that 3-furan carboxylic acid has
25 potent antimicrobial properties. Particularly surprising is the observation that this acid has a greater level of antimicrobial activity than 2-furan carboxylic acid. In contrast with the latter, 3-furan carboxylic acid has sufficient antimicrobial activity to be used as the sole antimicrobial in a cosmetic
30 formulation over a broad pH range. In particular, it is noted that 3-furan carboxylic acid is capable of preventing growth of Enterics, *Pseudomonas*, *Staphylococcus* and Mold, and substantially reducing the growth of yeast; in contrast, although 2-furan

carboxylic acid does exhibit inhibition of Enterics, *Pseudomonas*, and *Staphylococcus*, it has a very poor level of activity against Mold and yeast, making it inadequate on its own to preserve a cosmetic formulation. To achieve the desired antimicrobial effect, 3-furan carboxylic acid alone can be used in the formulation in an amount of from about 0.05-5%, preferably 0.1-3%, by weight of the total composition. Although the 2-furan carboxylic acid alone is insufficient protection, it may be desirable to use it in combination with 3-furan carboxylic acid as a supplement to that acid's activity, in an amount of 0.5-5%. In preferred formulations of the invention, both acids are combined, and in each case, throughout the specification and the claims, when the acid is referred to, it is understood that this refers to both the free acid and derivatives thereof having the same activity. These compounds are available commercially, for example, from Sigma Chemical.

It has also been observed that both 2- and 3-furan carboxylic acids possess a surprising level of anti-inflammatory activity. When tested in standard *in vitro* assays for anti-inflammatory activity, both compounds have been shown to be potent inhibitors of one or more steps in the inflammatory cascade. Both 2- and 3-furan carboxylic exhibit the ability to inhibit the adhesion of polymorphonuclear leukocytes (PMNs) to activated endothelial cells, an indicator of chronic, rather than acute, inflammation. In addition, both compounds are capable of inhibiting the activity of Phospholipase A₂ enzyme, which is a stimulator of arachidonic acid production; this is an indicator of inhibition of one of the early events in the inflammatory cascade of events, indicating an ability to inhibit acute inflammatory responses. In this particular property, however, 3-furan carboxylic acid is exceptionally active, inhibiting activity at a level of 0.01mg/ml. Therefore, in an additional embodiment of the invention, a cosmetic or pharmaceutical

composition for treatment or prevention of inflammation is provided, the composition comprising from anti-inflammatory effective amounts of 2-furan carboxylic acid, 3-furan carboxylic acid, or a mixture thereof. Generally speaking, the effective
5 amounts of these compounds will be in the same range as required for their antimicrobial activity. Such compositions are useful in the treatment and prevention of skin conditions or disorders which comprise an inflammatory aspect, for example, psoriasis, eczema, allergic contact dermatitis, or atopic dermatitis.

10 An additional useful component of the formulation is ferulic acid or derivatives thereof. Ferulic acid is also a naturally occurring material; it is found as a free acid in plants, and also occurs in ester form in seeds, leave and bark of plants. Esters are formed, for example, with long chain alcohols,
15 sterols, and hydroxyacids. Ferulic acid has previously been said to exhibit a variety of biological activities, such as antioxidant, deodorant, antiinflammatory, antimicrobial and anti-pruritic. It has not, however, to the inventors' knowledge, been identified as an antiirritant. In accordance with the
20 present invention, ferulic acid has been shown to have substantial antiirritant properties, which render it particularly useful in cosmetic formulations, and may be particularly desirable in combination with an active acid component that has the potential of causing irritation on the skin. The
25 antiirritant effect of ferulic acid can be achieved by application shortly prior to, or simultaneously with, or shortly after exposure of the skin to an irritating material. It may also be used alone to generally soothe sensitive, easily irritated skin. As used herein, the term "ferulic acid" will be
30 understood to encompass both the free acid and ester forms having the same activity as the free acid, unless otherwise specified. In the formulation, the ferulic acid is preferably used in an

amount of from about .05-2.5%, more preferably 0.1-1% by weight of the total composition.

It is particularly preferred that ferulic acid, or an ester thereof, is combined with both 2-furan carboxylic acid and 3-furan carboxylic acid, in the amounts specified above for each individual component. The formulations containing this combination exhibit sufficient antimicrobial control to avoid the use of conventional cosmetic or pharmaceutical preservatives. By "conventional" preservatives is meant those preservatives which have been routinely used in the past for inhibition of microbial growth; these include, but are not limited to, parabens, propionates, formaldehyde releasers, benzoates, and cresols. An added advantage of such formulations is that they are not only inherently non-irritating, but also reduce irritation already existing on the skin to which they are applied.

In an optional embodiment, the formulations may also comprise vanillin, vanillic acid, or vanillic acid esters, in an amount of from about .05-1%. Vanillin is also a naturally occurring material, which is known to have activity as an antimicrobial (Principles and Practice of Disinfection, Preservation and Sterilization, Russell et al, eds., Second Edition, Blackwell Scientific Publications; Cerrutti et al., J. Food Sci. 62: 608-610, 1997.)

For topical application, the active components can be formulated with a variety of cosmetically and/or pharmaceutically acceptable carriers. The term "pharmaceutically or cosmetically acceptable carrier" refers to a vehicle, for either pharmaceutical or cosmetic use, which vehicle delivers the active components to the intended target and which will not cause harm to humans or other recipient organisms. As used herein, "pharmaceutical" or "cosmetic" will be understood to encompass both human and animal pharmaceuticals or cosmetics. Useful carriers include, for example, water, acetone, ethanol, ethylene

glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, or mineral oil. Methodology and components for formulation of cosmetic and pharmaceutical compositions are well known, and can be found, for example, in Remington's

- 5 Pharmaceutical Sciences, Eighteenth Edition, A.R. Gennaro, Ed., Mack Publishing Co. Easton Pennsylvania, 1990. The carrier may be in any form appropriate to the mode of delivery, for example, solutions, colloidal dispersions, emulsions (oil-in-water or water-in-oil), suspensions, creams, lotions, gels, foams,
- 10 mousses, sprays and the like. The pH for the formulations of the invention can be from about 3 to about 8, preferably about 4-7, and most preferably about 4-6.

- The formulation, in addition to the carrier and the antimicrobial/antiirritant components, also can comprise other
- 15 optional materials which may be chosen depending on the carrier and/or the intended use of the formulation. Additional components include, but are not limited to, water soluble colorants (such as FD&C Blue #1); oil soluble colorants (such as D&C Green #6); water soluble sunscreens (such as Eusolex 232);
- 20 oil soluble sunscreens (such as Octyl Methoxycinnamate); particulate sunscreens (such as Zinc Oxide); antioxidants (such as BHT); chelating agents (such as Disodium EDTA); emulsion stabilizers (such as carbomer); preservatives (such as Methyl Paraben); fragrances (such as pinene); flavoring agents (such as
- 25 sorbitol); humectants (such as glycerine); waterproofing agents (such as PVP/Eicosene Copolymer); water soluble film-formers (such as Hydroxypropyl methylcellulose); oil-soluble film formers (such as Hydrogenated C-9 Resin); cationic polymers (such as Polyquaternium 10); anionic polymers (such as xanthan gum);
- 30 vitamins (such as Tocopherol); and the like.

The antimicrobial/antiirritant/antiinflammatory components are well-suited for combination with other active components intended for topical application. In particular, the ferulic

acid may assist in reducing the irritating effects of other active components in a formulation. Examples of known irritants that are frequently used for therapeutic purposes topically are retinoids, such as retinol and retinoic acid, and hydroxyacids, such as glycolic, lactic, or salicylic acids. Examples of other types of actives which may form part of the composition include, but are not limited to, those that improve or eradicate age spots, keratoses and wrinkles, analgesics, anesthetics, anti-acne agents, antibacterials, antiyeast agents, antifungal agents, antiviral agents, antidandruff agents, antidermatitis agents, antipruritic agents, antiemetics, antimotion sickness agents, anti-inflammatory agents, antihyperkeratolytic agents, anti-dry skin agents, antiperspirants, antipsoriatic agents, antiseborrheic agents, hair conditioners and hair treatment agents, antiaging agents, antiwrinkle agents, antiasthmatic agents and bronchodilators, sunscreen agents, antihistamine agents, skin lightening agents, depigmenting agents, wound-healing agents, vitamins, corticosteroids, tanning agents, sunscreens or hormones. More specific examples of useful active agents include retinoids, topical cardiovascular agents, clotrimazole, ketoconazole, miconazole, griseofulvin, hydroxyzine, diphenhydramine, pramoxine, lidocaine, procaine, mepivacaine, monobenzene, erythromycin, tetracycline, clindamycin, meclocyline, hydroquinone, minocycline, naproxen, ibuprofen, theophylline, cromolyn, albuterol, retinol, retinoic acid, 13-cis retinoic acid, hydrocortisone, hydrocortisone 21-acetate, hydrocortisone 17-valerate, hydrocortisone 17-butyrate, betamethasone valerate, betamethasone dipropionate, triamcinolone acetonide, fluocinonide, clobetasol, propionate, benzoyl peroxide, crotamiton, propranolol, promethazine, vitamin A palmitate, vitamin E acetate, DHEA and derivatives thereof, alpha- or beta-hydroxy acids, and mixtures thereof. The amount of

active agent to be used in any given formulation is readily determined in accordance with its usual dosage.

Particularly preferred embodiments of the present formulations are moisturizing creams or lotions. To that end, the antimicrobials/antiirritant components are combined with agents that are moisturizers, emollients or humectants. Examples of useful combinations are oils, fats, waxes, esters, fatty acid alcohols, fatty acid ethoxylates, glycols, sugars, hyaluronic acid and hyaluronates, dimethicone, cyclomethicone, and the like. Further examples can be found in the International Cosmetic Ingredient Dictionary, CTFA, Sixth Edition, 1995. The invention is further illustrated by the following non-limiting examples.

Application of the formulations of the invention is achieved in accordance with the nature and intended use of the final product. For example, a moisturizing, cleansing, or skin-soothing formulation may be used on a daily basis, or more or less frequently depending upon need. If the formulation contains a pharmaceutical or cosmetic active, the application will be in accordance with the recommended regimen for the active. Determination of other appropriate application regimens is a matter of routine optimization.

EXAMPLES

I. This example illustrates the efficacy of 3-furan carboxylic acid in retarding microbial growth in a cosmetic formulation.

A nonionic oil-in-water emulsion is prepared containing 0.473% 3-furan carboxylic acid. The formulation is divided into five portions, each one getting an inoculation at day 0, and a reinoculation after three weeks, of one of the following microbial cultures: (1)Enterics; (2)*Pseudomonas*; (3)*Staphylococcus*; (4)Yeast; and (5)Mold. The concentration of

microbes in each individual formulation is calculated on the day of the inoculation, and then again calculated on day 2, and at the end of one, two, three(pre-reinoculation) and four weeks, at a variety of pH values. These are also compared with formulations
5 containing 2-furan carboxylic acid and an identical series of microbes, and an acid pH control emulsion without a furan carboxylic acid added. The results are shown in Tables 1, 2 and 3. All numbers, other than those listed as "<10", are log base 10; thus, for example, "6.6" is equivalent to the presence of 8
10 million bacteria. Entries of <10 indicate less than 10 colony forming units, the lowest level of detection.

Table 1: 0.473% 3-furan carboxylic acidA. pH 3.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	<10	<10	<10	<10	<10
2	6.3	<10	<10	<10	<10	<10
3	6.3	<10	<10	<10	<10	<10
4	5.7	<10	<10	<10	<10	<10
5	5.2	<10	<10	<10	<10	<10

5 B. pH 4.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	<10	<10	<10	<10	<10
2	6.3	<10	<10	<10	<10	<10
3	6.3	<10	<10	<10	<10	<10
4	5.7	2.5	<10	<10	<10	<10
5	5.2	<10	<10	<10	<10	<10

C. pH 5.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	<10	<10	<10	<10	<10
2	6.3	<10	<10	<10	<10	<10
3	6.3	<10	<10	<10	<10	<10
4	5.7	2.5	2.7	1.7	2.0	1.8
5	5.2	<10	<10	<10	<10	<10

Table 2: 0.375% 2-furan carboxylic acidA. pH 3.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	<10	<10	<10	<10	<10
2	6.3	<10	<10	<10	<10	<10
3	6.3	<10	<10	<10	<10	<10
4	5.7	<10	<10	<10	<10	<10
5	5.2	<10	<10	<10	<10	<10

5

B. pH 4.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	<10	<10	<10	<10	<10
2	6.3	<10	<10	<10	<10	<10
3	6.3	<10	<10	<10	<10	<10
4	5.7	3.7	3.7	<10	<10	>5.7
5	5.2	1.6	<10	<10	<10	<10

C. pH 5.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	>6	>6	<10	<10	<10
2	6.3	>6	<10	<10	<10	<10
3	6.3	2.9	<10	<10	<10	<10
4	5.7	>5.7	>5.7	>5.7	>5.7	>5.7
5	5.2	5.2	5.2	5.2	3.0	>5

Table 3: Control emulsionA. pH 3.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	3.7	<10	<10	<10	<10
2	6.3	>6	<10	<10	<10	<10
3	6.3	2.0	<10	<10	<10	<10
4	5.7	5.7	5.7	5.7	4.5	>6
5	5.2	5.2	5.2	5.2	5.2	>5

B. pH 4.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	>6	>6	3.2	<10	>6
2	6.3	>6	>6	<10	<10	<10
3	6.3	>6	2.0	2.0	<10	<10
4	5.7	>6	>6	>6	>6	>6
5	5.2	5.2	5.2	5.2	3.3	>5

5.

C. pH 5.0

Pool #	Day 0	Day 2	Week 1	Week 2	Week 3	Week 4
1	6.6	>6	>6	>6	2.0	>6
2	6.3	>6	>6	<10	<10	<10
3	6.3	>6	>6	>6	4.2	<10
4	5.7	>6	>6	>6	>6	>6
5	5.2	>5.2	5.2	5.2	3.0	>6

10 Comparison of the antimicrobial effects of the two furan carboxylic acids shows that the 3-furan carboxylic acid is superior to 2-furan carboxylic acid in its ability to retard the growth of mold and yeast. The control shows that the antimicrobial effect of these two acids is due to the

characteristics of the specific acid, rather than the acid pH or the vehicle used.

Example 2: This example illustrates the antiirritant effect of
5 the components of the invention:

Seven volunteers with a history of skin sensitivity to Balsam of Peru are chosen for the study. The test compounds studied are as follows: Ferulic acid, 0.1% in hydroalcohol 1:1;
10 2-furan carboxylic acid, 0.1% in hydroalcohol 1:1; and 3-furan carboxylic acid, 0.1% in hydroalcohol, 1:1.

The test compounds are applied to the ventral forearms of panelists. The material is allowed to absorb for twenty minutes and then Balsam of Peru, an irritant is applied on the test
15 sites. Skin irritation is measured in terms of increase in skin redness. The degree of redness is measured with a Minolta Chromameter and compared with the controls, the positive control being skin treated with Balsam of Peru alone, and the negative control being a skin site treated with 10% cola solution(a known
20 antiirritant), and challenged like the test products.

Ferulic acid is 72% effective in preventing the onset of irritation. This compares favorably with the cola solution, which shows 69% reduction. Interestingly, both 2- and 3-furan carboxylic acid show some reduction in the onset of irritation,
25 at 49% and 48%, respectively.

Example 3: This example illustrates the anti-inflammatory effect of 2- and 3-furan carboxylic acids, in two different aspects of the inflammatory cascade.

30

A. PLA₂ (phospholipase A₂) inhibition

The enzyme reaction is accomplished in a cuvette containing 2m of phosphate buffered saline. Fluorometric substrate NBD-C6-HPC is

solubilized 1 mM in EtOH, and is diluted in the reaction mixture to yield a 10 μ M final concentration. The assay is initiated with the addition of .001 units per μ l of PLA₂ enzyme obtained from snake venom. Enzyme activity is monitored either with or without the presence of potential inhibitors. Materials tested are solubilized as follows: ferulic acid and 2-furan carboxylic acid are solubilized 10mg/ml in EtOH, and 3-furan carboxylic acid is solubilized 100mg/ml in EtOH; each sample is diluted appropriately to yield assay concentrations of .01, .1 and 1 mg/ml. Prior to enzyme addition, samples are measured fluorometrically at excitation 470 and emission 570 to determine background level of fluorescence. Following enzyme addition, samples are monitored every 15 minutes over a 60 minute period.

EC-50 values are calculated, but are limited to those samples which best yield a linear fit. Of those calculable in this manner, 2-furan carboxylic acid yielded a .5mg/ml value(Figure 1a). Although it was not possible to calculate an EC-50 for 3-furan carboxylic acid, as shown in Figure 1b, it provides the greatest inhibition by blocking the enzyme activity at .1mg/ml dose. Ferulic acid also shows some inhibitory effect, but is the weakest of the three tested(Figure 1c).

B. Inhibition of lymphocyte adhesion

Endothelial cells are grown to confluence and washed thoroughly with basal media prior to the experiment to remove all fetal calf serum. The test compounds are then added to the endothelial culture well and incubated for 2 hours. IL-1 β is added to stimulate expression of adhesion molecules on the endothelial cells before lymphocytes are added.

Lymphocytes are isolated from blood collected from adult human subjects and added to the endothelial culture wells. PHA is added to stimulate adhesion molecule expression in the lymphocytes and the cultures incubated overnight. After a brief

washing procedure, cells are incubated with Rose Bengal, and the number of adherent cells assessed by measuring the absorbance at 570nm.

Figure 2 illustrates that both 2- and 3-furan carboxylic acid are
5 very potent inhibitors of lymphocyte adhesion at concentrations
of 0.5% and 1%. The first bar shows the absorbance of
endothelial cells plus any adherent lymphocytes under non-
stimulated conditions. The second shows the increase in
absorbance after overnight incubation with PHA. The next four
10 bars show the decrease in absorbance back to non-stimulated
conditions in cultures preincubated with either 0.5% or 1% of the
two acids.

What we claim is:

1. A cosmetic or pharmaceutical formulation for topical application, the formulation comprising an antimicrobial effective amount of 3-furan carboxylic acid.
5
2. The formulation of claim 1 which also comprises an antimicrobial effective amount of 2-furan carboxylic acid.
- 10 3. The formulation of claim 1 which comprises an antiirritant effective amount of ferulic acid.
4. The formulation of claim 1 which comprises an antimicrobial effective amount of 2-furan carboxylic acid and an antiirritant effective amount of ferulic acid.
15
5. The formulation of claim 1 in which the acid is in free acid form.
- 20 6. The formulation of claim 1 in which each of the acids is in free acid form.
7. The formulation of claim 1 which also comprises vanillin.
- 25 8. The formulation of claim 4 which also comprises vanillin.
9. The formulation of claim 1 which is a moisturizing formulation.
- 30 10. The formulation of claim 4 which is a moisturizing formulation.

11. The formulation of claim 1 in which the amount of acid is from about 0.5-5%.
12. The formulation of claim 4 which comprises from 0.5-5% of each of free 2-furan carboxylic acid and 3-furan carboxylic acid, and from 0.5-2.5% of free ferulic acid.
13. A preservative-free cosmetic or pharmaceutical formulation comprising antimicrobial effective amounts of 3-furan carboxylic acid and 2-furan carboxylic acid, and an antiirritant effective amount of ferulic acid.
14. The formulation of claim 9 in which each of the acids is in free acid form.
15. A method for retarding microbial growth in a cosmetic or pharmaceutical formulation which comprises adding to the formulation an antimicrobial effective amount of 3-furan carboxylic acid.
16. A method for retarding microbial growth on skin which comprises applying to the skin an effective amount of a formulation of claim 1.
17. The method for retarding microbial growth on skin which comprises applying to the skin an effective amount of a formulation of claim 4.
18. The method for retarding microbial growth on skin which comprises applying to the skin an effective amount of a formulation of claim 12.

19. A method for preventing or reducing the irritation caused by an irritating material applied to the skin comprising applying to the skin an antiirritant effective amount of ferulic acid.
- 5 20. A cosmetic or pharmaceutical formulation for topical application comprising an antiirritant effective amount of ferulic acid.
- 10 21. The formulation of claim 20 which also comprises an irritating active component.
22. The formulation of claim 21 which also comprises an antimicrobial effective amount of 2-furan carboxylic acid and 3-furan carboxylic acid.
- 15 23. A method of inhibiting inflammation in the skin which comprises applying to the skin an effective amount of 2-furan carboxylic acid, 3-furan carboxylic acid, or a mixture thereof.
- 20 24. The method of claim 23 which comprises applying 3-furan carboxylic acid.
- 25 25. The method of claim 23 which comprises applying 2-furan carboxylic acid.
26. The method of claim 23 in which ferulic acid is also applied.
- 30 27. The method of claim 23 which comprises applying free 2-furan carboxylic acid, free 3-furan carboxylic acid, and free ferulic acid.

28. A method of treating a skin disorder having an inflammatory component comprising applying to the skin an effective amount of 2-furan carboxylic acid, 3-furan carboxylic acid, or a mixture thereof.

5

29. The method of claim 27 in which the skin disorder is selected from the group consisting of psoriasis, eczema, allergic contact dermatitis, or atopic dermatitis.


10 30. The method of claim 27 which comprises applying free 3-furan carboxylic acid.

31. The method of claim 27 which comprises applying free 2-furan carboxylic acid.


15

32. The method of claim 27 in which free ferulic acid is also applied.

20 33. The method of claim 27 which comprises applying free 2-furan carboxylic acid, free 3-furan carboxylic acid, and free ferulic acid.




(A)



(B)

FIGURE 1



(C)

FIGURE 1 (CONT'D)

3/3




FIG. 2

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.